



## The Crucial Role of Chloroplast-Related Proteins in Viral Genome Replication and Host Defense against Positive-Sense Single-Stranded RNA Viruses

John Bwalya<sup>1</sup> and Kook-Hyung Kim <sup>1,2,3\*</sup>

<sup>1</sup>Department of Agriculture Biotechnology, College of Agriculture and Life Sciences, Seoul National University, Seoul 08826, Korea

<sup>2</sup>Plant Genomics and Breeding Institute, Seoul National University, Seoul 08826, Korea

<sup>3</sup>Research of Institute Agriculture and Life Sciences, Seoul National University, Seoul 08826, Korea

(Received on October 4, 2022; Revised on December 19, 2022; Accepted on December 22, 2022)

**Plant viruses are responsible for worldwide production losses of numerous economically important crops. The most common plant RNA viruses are positive-sense single-stranded RNA viruses [(+ss RNA viruses)]. These viruses have small genomes that encode a limited number of proteins. The viruses depend on their host's machinery for the replication of their RNA genome, assembly, movement, and attraction to the vectors for dispersal. Recently researchers have reported that chloroplast proteins are crucial for replicating (+ss plant RNA viruses. Some chloroplast proteins, including translation initiation factor [eIF(iso)4E] and 75 DEAD-box RNA helicase RH8, help viruses fulfill their infection cycle in plants. In contrast, other chloroplast proteins such as PAP2.1, PSAC, and ATPsyn- $\alpha$  play active roles in plant defense against viruses. This is also consistent with the idea that reactive oxygen species, salicylic acid, jasmonic acid, and abscisic acid are produced in chloroplast. However, knowledge of molecular mechanisms and functions underlying these chloroplast host factors during the virus infection is still scarce and**

remains largely unknown. Our review briefly summarizes the latest knowledge regarding the possible role of chloroplast in plant virus replication, emphasizing chloroplast-related proteins. We have highlighted current advances regarding chloroplast-related proteins' role in replicating plant (+ss RNA viruses.

**Keywords :** chloroplast, chloroplast-virus interactions, plant defense, viral replication complex, virus replication

Viruses cause major crop losses worldwide and thus are a threat to sustainable and productive agriculture. New plant viruses are being discovered and continue to pose a clear danger to our food systems globally (Hilaire et al., 2022; Jones and Naidu, 2019; Rubio et al., 2020; Whitfield et al., 2015). Once positive-sense single-stranded RNA viruses [(+ss RNA viruses)] are inside the cell, the viral genome serves as a template for the production of a large number of progeny viruses (den Boon et al., 2010; Nagy and Pogany, 2008b), and RNA replication is done by viral proteins and host plant proteins in chloroplast membranes of the infected cell. Many viruses invade specific host chloroplast membranes in the process of viral genome replication (Hyodo and Okuno, 2016; Nagy et al., 2012; Sanfaçon, 2005; Xu and Nagy, 2014). Therefore, chloroplasts play central roles in replicating several plant virus species and biosynthesis of most plant hormones, making chloroplast factors crucial for plant defense response.

The chloroplast is a vital organelle of plant cells carrying out photosynthesis. In addition to the outer and inner membranes, mature chloroplasts have an internal membrane network of thylakoids, where the light energy is converted into chemical energy stored in ATP (Fig. 1)

\*Corresponding author.

Phone) +82-2-880-4677, FAX) +82-2-873-2317

E-mail) kookkim@snu.ac.kr

ORCID

Kook-Hyung Kim

<https://orcid.org/0000-0001-9066-6903>

Handling Editor : Seung-Kook Choi

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Articles can be freely viewed online at [www.pjonline.org](http://www.pjonline.org).

(Li et al., 2016). Many researchers have published quality data using confocal microscopes, and they have captured stunning and excellent images (Bhat et al., 2013; Kaido et al., 2014; Thivierge et al., 2008) which show that majority of the plant (+)ss RNA viruses replicate, assemble, and mature in chloroplast components called viral replication complexes (VRCs), including plum pox virus (PPV) (Martin et al., 1995), tobacco etch virus (Gadh and Hari, 1986), turnip mosaic virus (TuMV) (Kitajima and Costa, 1973), maize draft mosaic virus (Mayhew and Ford, 1974), and tobacco mosaic virus (TMV) (Bhat et al., 2013). Moreover, chloroplast-related proteins are reportedly involved in defense against plant viruses because chloroplasts are sites where defense-related hormones are synthesized (Bhattacharyya and Chakraborty, 2018; Bobik and Burch-Smith, 2015; Kozuleva et al., 2011; Nambara and Marion-Poll, 2005; Padmanabhan and Dinesh-Kumar, 2010; Seyfferth and Tsuda, 2014; Stael et al., 2015; Torres et al., 2006; Wasternack and Hause, 2013; Wildermuth et al., 2001; Yang et al., 2021). Recently, the chloroplast membrane-associated protein pSaC in photosystem I (PSI) and the ATP-synthase  $\alpha$ -subunit whose chloroplast DNA encoded protein is part of the ATP-synthase complex induced resistance in soybean mosaic virus (SMV) belonging to the *Potyvirus* genus (Bwalya et al., 2022).

In another example, infection with TMV reduced the expression levels of the *ATPsyn- $\gamma$*  gene and silencing *ATPsyn- $\gamma$*  in *Nicotiana benthamiana* increased TMV accumulation and pathogenicity (Bhat et al., 2013). Jiménez et al. (2006) have indicated a positive role for PSI-K protein (a product of the gene *psaK*) in resistance against the PPV. For PPV infection, transient expression of PSI-K decreased accumulation level of PPV RNA in *N. benthamiana*. But PPV RNA accumulation was enhanced when *psaK* was knocked down (Jiménez et al., 2006). Importantly, it should be noted that many viruses intrude photosynthetic machinery functionality through protein-protein interactions and participate in peddling these molecules to the chloroplast, where replication occurs (Fig. 2). This review summarizes the active roles of chloroplast proteins that play in the replication of (+)ss RNA viruses and plant defense against viruses.


### **Chloroplasts and Chloroplast-Related Proteins Facilitate the Viral Replication Cycle of Plant (+)ss RNA Viruses**

The viral replication process of (+)ss RNA viruses starts from the switch from translation to replication, including a selection of template RNA (Nagy and Pogany, 2006;

Nishikiori et al., 2006; Panavas et al., 2005). The viral and host components required for replication are targeted to chloroplast (subcellular) membranes, where the VRCs are assembled. VRCs synthesize the negative-sense (−) and (+) RNAs and release (+) RNA progenies (Fig. 1).

For plant (+)ss RNA viruses, host translation machinery uses the viral genome as an mRNA to produce replicating proteins and other viral proteins. Host and viral proteins have been well-documented to regulate the switch from translation to replication (Budziszewska and Obrepalska-Steplowska, 2018; Nagy and Pogany, 2008a). However, most host proteins are more likely to participate in the recruitment step. For example, the recruitment of brome mosaic virus RNAs for replication was affected by Lsm1p, Pat1p, and Dhh1p (Díez et al., 2000; Noueiry and Ahlquist, 2003). When the (+)ss RNA viruses come into the cell, the RNAs come out of the virions and are released into the cytoplasm. The genomes of (+)ss RNA viruses act as a template for translation to produce viral replication proteins, including RNA-dependent RNA polymerase (RdRp), using host machinery leading to a series of interactions between host translation factors and RNA replication (Dreher and Miller, 2006; Simon and Miller, 2013). The viral and host proteins are bound in a discriminatory manner to the (+)ss RNA template and target subcellular membranes. Translation and selection of the viral (+)ss RNA for replication takes place in the cytoplasm, whereas replication of (+)ss RNA viruses occurs on the surface of various intracellular membranes, including chloroplast membranes (Ahlquist, 2002; Ahlquist et al., 2003; Burgyan et al., 1996; den Boon et al., 2010; Rubino and Russo, 1998; Salonen et al., 2005; Widyasari et al., 2020). Intracellular targeting of the RdRp and viral (+)ss RNA to the replication site occurs in a favorable microenvironment to assemble VRCs. VRC formation at the replication site occurs through viral replicase, virus-encoded accessory proteins, and host factors recruited (Jin et al., 2017; Panavas et al., 2005; Salonen et al., 2005). The process of VRC formation is intimately associated with viral translation, intracellular movement, and intercellular movement. In early models of cell-to-cell movement, the viral MP alone was thought to be responsible for intracellular and intercellular movement. Proper coordination of these processes is required for efficient viral infection. Therefore, viruses must properly build VRCs to avoid or minimize disruption of a coordinated balance, in which host factors play a role in the assembly of viral replicase and regulate its function (Hafrén et al., 2010; Jungfleisch et al., 2015; Kaido et al., 2014).

The critical step is the synthesis of viral RNA (De Graaff



**Fig. 1.** Schematic representation of necessary steps for positive-sense single-stranded RNA [(+)ss RNA] virus genome replication. Following the entry into host cells, (1) viral genomic RNAs are released from virions into the cytoplasm, and (2) viral RNA [(+)ss RNA] translates RNA-dependent RNA polymerase (RdRp) at the early stage of infection and recruits additional factors (such as methyltransferase and chaperone). (3) The resulting viral replication proteins target themselves to recruit the host translation machinery for the successful production of viral replication proteins. (4) The recruited viral genomic (+) RNAs and host proteins are then trafficked to chloroplast membranes, where they assemble viral replication complexes (VRCs) on the host chloroplast membrane. VRCs are shown by an invagination of the plant chloroplast membrane containing the viral protein (blue shape), the viral RdRP (red shape), host proteins (green shape), and viral RNA (red line). The VRC synthesizes a complementary negative-strand RNA (green line) using the original (+)RNA as a template. The (−) RNA is then used as a template to synthesize many new (+)RNAs (red lines). Progeny viruses are released from VRCs, undergo additional translation and replication, or move to adjacent cells.

et al., 1993). The VRC synthesizes complementary (−) RNA using the original genomic (+) RNA as a template. The (−) RNA synthesizes more (+) RNAs. Newly synthesized (+) RNAs are then released from VRC to undergo rounds of translation and replication or move to adjacent cells or package into virions (Fig. 1). Nowadays, many studies have shown the association of the VRC with the outer chloroplast membrane (De Graaff et al., 1993; Moriceau et al., 2017; Nishikiori et al., 2006; Thivierge et

al., 2008). Table 1 summarizes chloroplast-related proteins in the viral replication process. It was reported a long time ago that VRCs of alfalfa mosaic virus (AMV), the genus *Alfamovirus*, are associated with the chloroplast outer membrane (De Graaff et al., 1993). Turnip yellow mosaic virus (TYMV; a (+)ss RNA virus that shares replication features with other alphavirus-like supergroups) replication also occurs in close association with the chloroplast outer envelope membranes (Hatta et al., 1973; Koonin and Dolja,

**Table 1.** Chloroplast factors and their cellular localization during virus replication

Plant virus	Chloroplast factor	Subcellular localization	Reference
Soybean mosaic virus (SMV)	GmPAP2.1	Chloroplast	Widyasari et al. (2022)
	Rieske Fe/S	Chloroplast membrane	Shi et al. (2007)
	GmATPsy- $\alpha$	Chloroplast membrane, cytoplasm, and nucleolus	Bwalya et al. (2022)
	GmPSaC	Chloroplast membrane, cytoplasm, and nucleolus	Bwalya et al. (2022)
Bamboo mosaic virus	NbLTP1	Chloroplast	Shi et al. (2007)
	cPGK	Chloroplast and cytoplasm	Cheng et al. (2013)
Tobacco vein-mottling virus (TVMV)	PSI-K	Chloroplast membrane	Jiménez et al. (2006)
Tobacco vein banding mosaic virus (TVBMV)	NbRPL1	Chloroplast	Cheng et al. (2021)
Alfalfa mosaic virus (AMV)	PsbP	Cytoplasm	Balasubramaniam et al. (2014)
Cucumber mosaic virus Y strain	ChII mRNA	Cytoplasm	Shimura et al. (2011)
Potato mop-top virus	Chloroplast lipid	Chloroplast membrane	Cowan et al. (2012)
Potato virus X	Plastocyanin	Chloroplast	Qiao et al. (2009)
Alternanthera mosaic virus (AltMV)	PsbO	Chloroplast membrane	Jang et al. (2013)

1993; Prod'homme et al., 2001). The TYMV 140K protein was previously shown to be responsible for the recruitment of the polymerase to VRCs and targeting the VRCs to the outer chloroplast membrane where viral replication occurs (Jakubiec et al., 2004; Prod'homme et al., 2003). Moriceau et al. (2017) investigated determinants for the *in vivo* chloroplast targeting of the TYMV 140K replication protein. They identified the two amphipathic helices  $\alpha$ A and  $\alpha$ B within 140K/98K that constitute the determinants for chloroplast targeting of the TYMV VRCs. However, detailed delivery mechanisms remain to be determined.

Furthermore, published data for 3D electron tomography of barley stripe mosaic virus revealed replication factories and remodeling of the chloroplast outer membranes, characterized by clustering outer membrane-invaginated spherules in inner membrane-derived packets (Budziszewska and Obrepalska-Steplowska, 2018). Our recent study also showed that chloroplast-related proteins pSaC and ATPsyn- $\alpha$  were localized in the chloroplast envelope (Bwalya et al., 2022). Nonetheless, helicases and chloroplast factors participate in each step of RNA synthesis. One such example of (+)ss RNA viruses infecting plant species is the eukaryotic translation elongation factor 1A (eEF1A), which was found to be a permanent resident of the tombusvirus replicase complex and eEF1A promotes the synthesis of (-) RNA by replicase complex (Li et al., 2009). The elevated abundance of eEF1A in the cells and eEF1A interaction with viral (+) RNAs, including the 3' untranslated region of TYMV

(Dreher, 1999), TMV, and TuMV, might facilitate the recruitment of eEF1A into VRC (Nishikiori et al., 2006; Thivierge et al., 2008). In TuMV infection, 6K2 vesicles were transported to the chloroplast and accumulated at the chloroplast membrane, where they induce the formation of chloroplast-bound elongated tubular structures and ultimately form chloroplast aggregation (Wei et al., 2013). Plant viruses co-opt host proteins such as methyltransferase and chaperone for viral replication (Budziszewska and Obrepalska-Steplowska, 2018), which may perform similar tasks in viral replication. During viral infection, an RNA helicase protein, ISE2 (increased size exclusion limit 2), was highly expressed and accumulated in the chloroplast affecting viral replication and cell-to-cell communication (Ganusova et al., 2017).

### Chloroplast-Related Proteins Interact with Viral Components during Virus Replication

The accumulating evidence highlights that chloroplasts and chloroplast-related proteins can interact with viral components to favor the replication and movement of (+) ss RNA viruses (Bobik and Burch-Smith, 2015; Zhao et al., 2016, 2019) (Table 2, Fig. 2). In soybean (*Glycine max*), purple acid phosphatase (GmPAP2.1) conferred salicylic acid (SA)-dependent resistance to a susceptible cultivar by interacting with the SMV P1 protein in the chloroplasts, transient knockdown of endogenous SA-related genes resulted in systemic infection by SMV

**Table 2.** Summary of viral protein-chloroplast factor interactions

Plant virus	Virus component	Chloroplast factor	Reference
Soybean mosaic virus (SMV)	P1	GmATPsy- $\alpha$	Bwalya et al. (2022)
		GmPAP2.1	Widyasari et al. (2022)
		GmPSaC	Bwalya et al. (2022)
		Rieske Fe/S	Shi et al. (2007)
Plum pox virus	CI	PSI-K	Jiménez et al. (2006), Martin et al. (1995)
Tobacco vein-mottling virus (TVMV)	CI	PsaK	Caplan et al. (2008)
		PSI-K	Jiménez et al. (2006)
Tobacco mosaic virus	RNA replicase	PsbO	Abbink et al. (2002)
	126K replicase	NRIP 1	Caplan et al. (2008)
	126K/183K replicase	ATP synthase- $\gamma$ subunit (AtpC) Rubisco activase (RCA)	Bhat et al. (2013) Bhat et al. (2013)
Tomato mosaic virus	MP	RbcS	Zhao et al. (2013)
Potato virus Y	HC-Pro	MinD	Jin et al. (2007)
Alfalfa mosaic virus (AMV)	CP	PsbP	Balasubramaniam et al. (2014)
Sugarcane mosaic virus	HC-Pro	Ferredoxin	Cheng et al. (2008)
Turnip mosaic virus	VPg	NADH	Zhai et al. (2021)
	P3	Rubisco	Lin et al. (2011)
Alternanthera mosaic virus (AltMV)	TGB3	PsbO	Jang et al. (2013)


strain G5H (Widyasari et al., 2022). In *N. benthamiana*, overexpression of chloroplast nicotinamide adenine dinucleotide dehydrogenase-like complex M subunit gene (*NdhM*) inhibited TuMV accumulation, and the localization of NbNdhM is altered by its interaction with TuMV VPg in a way that promoted virus infection (Zhai et al., 2021). Cylindrical inclusion (CI) protein from PPV interacted with the PSI PSI-K protein (a product of the gene *psaK*) of *N. benthamiana*. Transient coexpression of PPV CI in *N. benthamiana* leaves decreased the level of PSI-K while silencing of *psaK* enhanced PPV accumulation (Jiménez et al., 2006). A chloroplast ribosomal protein large subunit 1 (NbRPL1), localized to the chloroplasts via its transit peptide, interacted with tobacco vein banding mosaic virus (TVBMV) nuclear inclusion protein b (NIb) and enhanced TVBMV infection. Silencing of NbRPL1 expression reduced TVBMV replication in *N. benthamiana* (Cheng et al., 2021). The photosystem II (PSII) oxygen-evolving complex protein (PsbP) interacted with the AMV capsid protein, and its overexpression reduced virus replication (Balasubramaniam et al., 2014). The helicase domain of the TMV replicase interacted with the *psbO*-encoded 33-kDa protein, a component of the oxygen-evolving complex. TMV infection depleted the *psbO* gene and the thylakoid's entire PSII core complex. Silencing of the *psbO* gene

increased TMV replication (Abbink et al., 2002; Balasubramaniam et al., 2014; Lehto et al., 2003). In maize (*Zea mays*), the multifunctional helper component-proteinase protein of sugar cane mosaic virus (SCMV) interacted with chloroplast precursor of ferredoxin-5 (FdV), and SCMV infection significantly downregulated the expression level of *FdV* mRNA in maize plants (Cheng et al., 2008). SMV P1 protein interacted with Rieske Fe/S protein in tobacco, retarded the Fe/S transport to the chloroplast (Shi et al., 2007). Many more interactions between chloroplast and viral proteins are listed in Table 2.

### Chloroplast Plays a Vital Role in Plant Antiviral Defense

The chloroplast is one of the crucial organelles that provides energy and carbon through photosynthesis. However, the plant defense mechanism tends to decrease the photosynthetic rate and other anabolic processes to reduce the organic carbon supply to pathogens (Serrano et al., 2016).

Other than photosynthesis, chloroplast plays significant roles in plant defense. It provides plant defense against viruses because of the production of secondary metabolites, including calcium, reactive oxygen species (ROS), and



**Fig. 2.** Schematic representation of the important events during chloroplast host factors and potyvirus interactions in a plant cell. After entry into host plant cells, a potyvirus virion undergoes the disassembly of viral particles and releases the viral genome. Viral genomic RNA is then used as the template for translation to produce viral polyproteins (11 viral proteins indicated by grey squares). The 6K2 remodels the subcellular membranes to form the viral replication complexe (VRC)-containing vesicles for potyvirus genome replication. The 6K2-induced vesicles may subsequently target chloroplasts for robust viral replication. The NIb is recruited to the VRC, likely via its interaction with the VPg domain of 6K2–VPg-Pro. Then, NIb recruits many host factors, such as eEF1A, PSaC, and ATP-synthase  $\alpha$ -subunit. In the figure, each viral protein is represented by a grey color-coded square in a location where they play a crucial role. A light blue color-coded semi-circle indicates chloroplast-related host proteins identified for virus infection. They are depicted in sites where they interact with viral protein or play a crucial role.

biosynthesis of several defense-related hormones like SA, jasmonic acid (JA), and abscisic acid (ABA) that have an important connection with plant immunity (Bobik and Burch-Smith, 2015; Kozuleva et al., 2011; Nambara and Marion-Poll, 2005; Padmanabhan and Dinesh-Kumar, 2010; Seyfferth and Tsuda, 2014; Stael et al., 2015; Torres et al., 2006; Wasternack and Hause, 2013; Wildermuth et al., 2001).

Most plant virus-induced SA is synthesized through the isochorismate pathway in chloroplast and plays crucial

roles in plant defense against viruses, and is essential for local and systemic acquired resistance (Alazem and Lin, 2015; Boatwright and Pajerowska-Mukhtar, 2013; Seyfferth and Tsuda, 2014; Zhao et al., 2016). For example, overexpression of GmPAP2.1, a chloroplast-localized protein, resulted in the upregulation of the SA pathway. Overexpression of GmPAP2.1 showed resistance to SMV infection, while transient knockdown of endogenous SA-related genes caused severe systemic symptoms by SMV (Widyasari et al., 2022). Another chloroplast-localized

protein, named calcium-sensing receptor, acts upstream of SA accumulation and connects chloroplasts to cytoplasmic-nuclear immune responses in *Arabidopsis thaliana* (Nomura et al., 2012).

JA is synthesized from linolenic acid by the octadecanoid pathway, and biosynthesis starts with the conversion of linolenic acid to 12-oxo-phytodienoic acid in the chloroplast membranes (Turner et al., 2002). JA also plays a crucial role in plant-virus interaction. For example, in silencing the JA perception gene, *COII* (coronatine insensitive 1) accelerates the development of symptoms caused by the co-infection of potato virus X (PVX) and potato virus Y. It increases the level of viral titers at the early stages of infection (García-Marcos et al., 2013).

Pathways of ABA are also involved in plant resistance to viruses (Alazem and Lin, 2015; Alazem et al., 2017, 2018, 2019; Nomura et al., 2012; Zhang et al., 2012; Zhao et al., 2016). Previously published reports showed that ABA enhanced the expression of the antiviral RNA silencing genes in soybean and *A. thaliana* and that the enhanced expression confers partial resistance against SMV, bamboo mosaic virus, and PVX (Alazem et al., 2017, 2019).

The reaction centers of PSI and PSII in chloroplast thylakoids are the primary generation site of ROS. ROS is a sign of activated antiviral defense (Bwalya et al., 2022; Calil and Fontes, 2017; Wu et al., 2017). The reaction centers of PSI and PSII in chloroplast thylakoids produce ROS and the photosynthetic electron transport chain. Superoxide anion ( $O_2^-$ ) is the primary reduced product of  $O_2$  photoreduction, and its disproportionation produces  $H_2O_2$  in chloroplast thylakoids membranes (Asada, 2006; Mühlenbock et al., 2008; Zhao et al., 2016). ROS and calcium bursts in chloroplasts activate signaling cascades that regulate the expression of defense-related genes; therefore, ROS and calcium bursts act as chloroplast-to-nucleus retrograde signals when plants recognize the early step of virus infection (Bhattacharyya and Chakraborty, 2018; Medina-Puche et al., 2020; Nomura et al., 2012; Zhao et al., 2016).

The burst of intracellular ROS can be detected during virus infection in incompatible and compatible interactions (Allan et al., 2001; Hakmaoui et al., 2012). Chloroplast-sourced ROS are essential for hypersensitive response induced by the incompatible defensive response (Torres et al., 2006; Zurbriggen et al., 2010).

## Conclusions and Future Perspectives

Chloroplasts have been recognized as a common target by many plant viruses. Hence chloroplast-virus interaction is

an epicenter of plant-virus interplays. Viruses may directly modify chloroplast membranes to assemble their replication complex for viral genome replication. Based on previously published reports (Balasubramaniam et al., 2014; Bhat et al., 2013; Bhattacharyya and Chakraborty, 2018; Budziszewska and Obrepalska-Steplowska, 2018; Bwalya et al., 2022; Cheng et al., 2021; Jiménez et al., 2006; Nagy et al., 2012; Serrano et al., 2016; Wei et al., 2013; Widyasari et al., 2022; Yang et al., 2021), we can conclude that some chloroplast-related proteins could function in virus replication, while some are involved in inhibiting viruses. Our review has summarized a few chloroplast-related proteins identified by researchers and their possible roles in virus infection. However, more chloroplast-related proteins must be determined to understand the proteins involved in host defense accurately. If the desired chloroplast-related gene functions are observed from host plants, they might be used to genetically engineer other plants to express these gene products after isolation and cloning. For example, scientists have boosted a carbon-craving enzyme called RuBisCO to turbocharge photosynthesis in corn (Salesse-Smith et al., 2018). The discovery promises to be a critical step in improving agricultural efficiency and yield. Moreover, more reports showed that overexpression of photosynthesis genes could enhance virus resistance in maize and soybean plants (Bwalya et al., 2022; Wang et al., 2018; Widyasari et al., 2022).

## Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

## Acknowledgments

This research was supported by grants from the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, and Forestry (120080-05-1-HD030), funded by the Ministry of Agriculture, Food and Rural Affairs, Republic of Korea. JB was supported by a research fellowship from the Brain Korea 21 Four Program.

## References

- Abbink, T. E. M., Peart, J. R., Mos, T. N. M., Baulcombe, D. C., Bol, J. F. and Linthorst, H. J. M. 2002. Silencing of a gene encoding a protein component of the oxygen-evolving complex of photosystem II enhances virus replication in plants. *Virology* 295:307-319.
- Ahlquist, P. 2002. RNA-dependent RNA polymerases, viruses, and RNA silencing. *Science* 296:1270-1273.

- Ahlquist, P., Noueiry, A. O., Lee, W.-M., Kushner, D. B. and Dye, B. T. 2003. Host factors in positive-strand RNA virus genome replication. *J. Virol.* 77:8181-8186.
- Alazem, M., He, M.-H., Moffett, P. and Lin, N.-S. 2017. Abscisic acid induces resistance against bamboo mosaic virus through Argonaute 2 and 3. *Plant Physiol.* 174:339-355.
- Alazem, M., Kim, K.-H. and Lin, N.-S. 2019. Effects of abscisic acid and salicylic acid on gene expression in the antiviral RNA silencing pathway in *Arabidopsis*. *Int. J. Mol. Sci.* 20:2538.
- Alazem, M. and Lin, N.-S. 2015. Roles of plant hormones in the regulation of host-virus interactions. *Mol. Plant Pathol.* 16:529-540.
- Alazem, M., Tseng, K.-C., Chang, W.-C., Seo, J.-K. and Kim, K.-H. 2018. Elements involved in the Rsv3-mediated extreme resistance against an avirulent strain of soybean mosaic virus. *Viruses* 10:581.
- Allan, A. C., Lapidot, M., Culver, J. N. and Fluhr, R. 2001. An early tobacco mosaic virus-induced oxidative burst in tobacco indicates extracellular perception of the virus coat protein. *Plant Physiol.* 126:97-108.
- Asada, K. 2006. Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol.* 141:391-396.
- Balasubramaniam, M., Kim, B.-S., Hutchens-Williams, H. M. and Loesch-Fries, L. S. 2014. The photosystem II oxygen-evolving complex protein PsbP interacts with the coat protein of alfalfa mosaic virus and inhibits virus replication. *Mol. Plant-Microbe Interact.* 27:1107-1118.
- Bhat, S., Folimonova, S. Y., Cole, A. B., Ballard, K. D., Lei, Z., Watson, B. S., Sumner, L. W. and Nelson R. S. 2013. Influence of host chloroplast proteins on tobacco mosaic virus accumulation and intercellular movement. *Plant Physiol.* 161:134-147.
- Bhattacharyya, D. and Chakraborty, S. 2018. Chloroplast: the Trojan horse in plant–virus interaction. *Mol. Plant Pathol.* 19:504-518.
- Boatwright, J. L. and Pajerowska-Mukhtar, K. 2013. Salicylic acid: an old hormone up to new tricks. *Mol. Plant Pathol.* 14:623-634.
- Bobik, K. and Burch-Smith, T. M. 2015. Chloroplast signaling within, between and beyond cells. *Front. Plant Sci.* 6:781.
- Budziszewska, M. and Obrepalska-Steplowska, A. 2018. The role of the chloroplast in the replication of positive-sense single-stranded plant RNA viruses. *Front. Plant Sci.* 9:1776.
- Burgyan, J., Rubino, L. and Russo, M. 1996. The 5'-terminal region of a tombusvirus genome determines the origin of multi-vesicular bodies. *J. Gen. Virol.* 77:1967-1974.
- Bwalya, J., Alazem, M. and Kim, K.-H. 2022. Photosynthesis-related genes induce resistance against soybean mosaic virus: evidence for involvement of the RNA silencing pathway. *Mol. Plant Pathol.* 23:543-560.
- Calil, I. P. and Fontes, E. P. B. 2017. Plant immunity against viruses: antiviral immune receptors in focus. *Ann. Bot.* 119:711-723.
- Caplan, J. L., Mamillapalli, P., Burch-Smith, T. M., Czymmek, K. and Dinesh-Kumar, S. P. 2008. Chloroplastic protein NRIP1 mediates innate immune receptor recognition of a viral effector. *Cell* 132:449-462.
- Cheng, D.-J., Xu, X.-J., Yan, Z.-Y., Tettey, C. K., Fang, L., Yang, G.-L., Geng, C., Tian Y.-P. and Li, X.-D. 2021. The chloroplast ribosomal protein large subunit 1 interacts with viral polymerase and promotes virus infection. *Plant Physiol.* 187:174-186.
- Cheng, S.-F., Huang, Y.-P., Chen, L.-H., Hsu, Y.-H. and Tsai, C.-H. 2013. Chloroplast phosphoglycerate kinase is involved in the targeting of bamboo mosaic virus to chloroplasts in *Nicotiana benthamiana* plants. *Plant Physiol.* 163:1598-1608.
- Cheng, Y.-Q., Liu, Z.-M., Xu, J., Zhou, T., Wang, M., Chen, Y.-T., Li, H.-F. and Fan, Z.-F. 2008. HC-Pro protein of sugar cane mosaic virus interacts specifically with maize ferredoxin-5 *in vitro* and *in planta*. *J. Gen. Virol.* 89:2046-2054.
- Cowan, G. H., Roberts, A. G., Chapman, S. N., Ziegler, A., Savenkov, E. I. and Torrance, L. 2012. The potato mop-top virus TGB2 protein and viral RNA associate with chloroplasts and viral infection induces inclusions in the plastids. *Front. Plant Sci.* 3:290.
- De Graaff, M., Coscoy, L. and Jaspars, E. M. J. 1993. Localization and biochemical characterization of alfalfa mosaic virus replication complexes. *Virology* 194:878-881.
- den Boon, J. A., Diaz, A. and Ahlquist, P. 2010. Cytoplasmic viral replication complexes. *Cell Host Microbe* 8:77-85.
- Díez, J., Ishikawa, M., Kaido, M. and Ahlquist, P. 2000. Identification and characterization of a host protein required for efficient template selection in viral RNA replication. *Proc. Natl. Acad. Sci. U. S. A.* 97:3913-3918.
- Dreher, T. W. 1999. Functions of the 3'-untranslated regions of positive strand RNA viral genomes. *Annu. Rev. Phytopathol.* 37:151-174.
- Dreher, T. W. and Miller, W. A. 2006. Translational control in positive strand RNA plant viruses. *Virology* 344:185-197.
- Gadh, I. P. S. and Hari, V. 1986. Association of tobacco etch virus related RNA with chloroplasts in extracts of infected plants. *Virology* 150:304-307.
- Ganusova, E. E., Rice, J. H., Carlew, T. S., Patel, A., Perrodin-Njoku, E., Hewezi, T. and Burch-Smith, T. M. 2017. Altered expression of a chloroplast protein affects the outcome of virus and nematode infection. *Mol. Plant-Microbe Interact.* 30:478-488.
- García-Marcos, A., Pacheco, R., Manzano, A., Aguilar, E. and Tenllado, F. 2013. Oxylipin biosynthesis genes positively regulate programmed cell death during compatible infections with the synergistic pair potato virus X–potato virus Y and tomato spotted wilt virus. *Virol. J.* 87:5769-5783.
- Hafrén, A., Hofius, D., Rönnholm, G., Sonnewald, U. and Mäkinen, K. 2010. HSP70 and its cochaperone CPIP promote potyvirus infection in *Nicotiana benthamiana* by regulating viral coat protein functions. *Plant Cell* 22:523-535.

- Hakmaoui, A., Pérez-Bueno, M. L., García-Fontana, B., Camejo, D., Jiménez, A., Sevilla, F. and Barón, M. 2012. Analysis of the antioxidant response of *Nicotiana benthamiana* to infection with two strains of pepper mild mottle virus. *J. Exp. Bot.* 63:5487-5496.
- Hatta, T., Bullivant, S. and Matthews, R. E. 1973. Fine structure of vesicles induced in chloroplasts of Chinese cabbage leaves by infection with turnip yellow mosaic virus. *J. Gen. Virol.* 20:37-50.
- Hilaire, J., Tindale, S., Jones, G., Pingarron-Cardenas, G., Bačnik, K., Ojo, M. and Frewer, L. J. 2022. Risk perception associated with an emerging agri-food risk in Europe: plant viruses in agriculture. *Agric. Food Secur.* 11:21.
- Hyodo, K. and Okuno, T. 2016. Pathogenesis mediated by proviral host factors involved in translation and replication of plant positive-strand RNA viruses. *Curr. Opin. Virol.* 17:11-18.
- Jakubiec, A., Notaise, J., Tournier, V., Héricourt, F., Block, M. A., Drugeon, G., van Aelst, L. and Jupin, I. 2004. Assembly of turnip yellow mosaic virus replication complexes: interaction between the proteinase and polymerase domains of the replication proteins. *J. Virol.* 78:7945-7957.
- Jang, C., Seo, E.-Y., Nam, J., Bae, H., Gim, Y. G., Kim, H. G., Cho, I. S., Lee, Z.-W., Bauchan, G. R., Hammond, J. and Lim, H.-S. 2013. Insights into alternanthera mosaic virus TGB3 functions: interactions with *Nicotiana benthamiana* PsbO correlate with chloroplast vesiculation and veinal necrosis caused by TGB3 over-expression. *Front. Plant Sci.* 4:5.
- Jiménez, I., López, L., Alamillo, J. M., Valli, A. and García, J. A. 2006. Identification of a plum pox virus CI-interacting protein from chloroplast that has a negative effect in virus infection. *Mol. Plant-Microbe Interact.* 19:350-358.
- Jin, X., Jiang, Z., Zhang, K., Wang, P., Cao, X., Yue, N., Wang, X., Zhang, X., Li, Y., Li, D., Kang, B.-H. and Zhang, Y. 2017. Three-dimensional analysis of chloroplast structures associated with virus infection. *Plant Physiol.* 176:282-294.
- Jin, Y., Ma, D., Dong, J., Li, D., Deng, C., Jin, J. and Wang, T. 2007. The HC-pro protein of potato virus Y interacts with NtMinD of tobacco. *Mol. Plant-Microbe Interact.* 20:1505-1511.
- Jones, R. A. C. and Naidu, R. A. 2019. Global dimensions of plant virus diseases: current status and future perspectives. *Annu. Rev. Virol.* 6:387-409.
- Jungfleisch, J., Chowdhury, A., Alves-Rodrigues, I., Tharun, S. and Díez, J. 2015. The Lsm1-7-Pat1 complex promotes viral RNA translation and replication by differential mechanisms. *RNA* 21:1469-1479.
- Kaido, M., Abe, K., Mine, A., Hyodo, K., Taniguchi, T., Taniguchi, H., Mise, K. and Okuno, T. 2014. GAPDH: a recruits a plant virus movement protein to cortical virus replication complexes to facilitate viral cell-to-cell movement. *PLoS Pathog.* 10:e1004505.
- Kitajima, E. W. and Costa, A. S. 1973. Aggregates of chloroplasts in local lesions induced in *Chenopodium quinoa* Wild. by turnip mosaic virus. *J. Gen. Virol.* 20:413-416.
- Koonin, E. V. and Dolja, V. V. 1993. Evolution and taxonomy of positive-strand RNA viruses: implications of comparative analysis of amino acid sequences. *Crit. Rev. Biochem. Mol. Biol.* 28:375-430.
- Kozuleva, M., Klenina, I., Proskuryakov, I., Kirilyuk, I. and Ivanov, B. 2011. Production of superoxide in chloroplast thylakoid membranes: ESR study with cyclic hydroxylamines of different lipophilicity. *FEBS Lett.* 585:1067-1071.
- Lehto, K., Tikkannen, M., Hiriart, J.-B., Paakkarinen, V. and Aro, E.-M. 2003. Depletion of the photosystem II core complex in mature tobacco leaves infected by the flavum strain of tobacco mosaic virus. *Mol. Plant-Microbe Interact.* 16:1135-1144.
- Li, Y., Cui, H., Cui, X. and Wang, A. 2016. The altered photosynthetic machinery during compatible virus infection. *Curr. Opin. Virol.* 17:19-24.
- Li, Z., Pogany, J., Panavas, T., Xu, K., Esposito, A. M., Kinzy, T. G. and Nagy P. D. 2009. Translation elongation factor 1A is a component of the tombusvirus replicase complex and affects the stability of the p33 replication co-factor. *Virology* 385:245-260.
- Lin, L., Luo, Z., Yan, F., Lu, Y., Zheng, H. and Chen, J. 2011. Interaction between potyvirus P3 and ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO) of host plants. *Virus Genes* 43:90-92.
- Martin, M. T., Cervera, M. T., Garcia, J. A. and Bonay, P. 1995. Properties of the active plum pox potyvirus RNA polymerase complex in defined glycerol gradient fractions. *Virus Res.* 37:127-137.
- Mayhew, D. E. and Ford, R. E. 1974. Detection of ribonuclease-resistant RNA in chloroplasts of corn leaf tissue infected with maize dwarf mosaic virus. *Virology* 57:503-509.
- Medina-Puche, L., Tan, H., Dogra, V., Wu, M., Rosas-Diaz, T., Wang, L., Ding, X., Zhang, D., Fu, X., Kim, C. and Lozano-Duran, R. 2020. A defense pathway linking plasma membrane and chloroplasts and co-opted by pathogens. *Cell* 182:1109-1124.
- Moriceau, L., Jomat, L., Bressanelli, S., Alcaide-Lordan, C. and Jupin, I. 2017. Identification and molecular characterization of the chloroplast targeting domain of turnip yellow mosaic virus replication proteins. *Front. Plant Sci.* 8:2138.
- Mühlenbock, P., Szechynska-Hebda, M., Plaszczycy, M., Baudo, M., Mateo, A., Mullineaux, P. M., Parker, J. E., Karpinska, B. and Karpinski, S. 2008. Chloroplast signaling and LESSION SIMULATING DISEASE1 regulate crosstalk between light acclimation and immunity in *Arabidopsis*. *Plant Cell* 20:2339-2356.
- Nagy, P. D., Barajas, D. and Pogany, J. 2012. Host factors with regulatory roles in tombusvirus replication. *Curr. Opin. Virol.* 2:691-698.
- Nagy, P. D. and Pogany, J. 2006. Yeast as a model host to dissect functions of viral and host factors in tombusvirus replication. *Virology* 344:211-220.
- Nagy, P. D. and Pogany, J. 2008a. Host factors promoting viral

- RNA replication. *Viral Genome Replication* 1:267-295.
- Nagy, P. D. and Pogany, J. 2008b. Multiple roles of viral replication proteins in plant RNA virus replication. *Methods Mol. Biol.* 451:55-68.
- Nambara, E. and Marion-Poll, A. 2005. Abscisic acid biosynthesis and catabolism. *Annu. Rev. Plant Biol.* 56:165-185.
- Nishikiori, M., Dohi, K., Mori, M., Meshi, T., Naito, S. and Ishikawa, M. 2006. Membrane-bound tomato mosaic virus replication proteins participate in RNA synthesis and are associated with host proteins in a pattern distinct from those that are not membrane bound. *J. Virol.* 80:8459-8468.
- Nomura, H., Komori, T., Uemura, S., Kanda, Y., Shimotani, K., Nakai, K., Furuichi, T., Takebayashi, K., Sugimoto, T., Sano, S., Suwastika, I. N., Fukusaki, E., Yoshioka, H., Nakahira, Y. and Shiina, T. 2012. Chloroplast-mediated activation of plant immune signalling in *Arabidopsis*. *Nat. Commun.* 3:926.
- Noueiry, A. O. and Ahlquist, P. 2003. Brome mosaic virus RNA replication: revealing the role of the host in RNA virus replication. *Annu. Rev. Phytopathol.* 41:77-98.
- Padmanabhan, M. S. and Dinesh-Kumar, S. P. 2010. All hands on deck: the role of chloroplasts, endoplasmic reticulum, and the nucleus in driving plant innate immunity. *Mol. Plant-Microbe Interact.* 23:1368-1380.
- Panavas, T., Hawkins, C. M., Panaviene, Z. and Nagy, P. D. 2005. The role of the p33:p33/p92 interaction domain in RNA replication and intracellular localization of p33 and p92 proteins of cucumber necrosis tombusvirus. *Virology* 338:81-95.
- Prod'homme, D., Jakubiec, A., Tournier, V., Drugeon, G. and Jupin, I. 2003. Targeting of the turnip yellow mosaic virus 66K replication protein to the chloroplast envelope is mediated by the 140K protein. *J. Virol.* 77:9124-9135.
- Prod'homme, D., Le Panse, S., Drugeon, G. and Jupin, I. 2001. Detection and subcellular localization of the turnip yellow mosaic virus 66K replication protein in infected cells. *Virology* 281:88-101.
- Qiao, Y., Li, H. F., Wong, S. M. and Fan, Z. F. 2009. Plastocyanin transit peptide interacts with potato virus X coat protein, while silencing of plastocyanin reduces coat protein accumulation in chloroplasts and symptom severity in host plants. *Mol. Plant-Microbe Interact.* 22:1523-1534.
- Rubio, L., Galipienso, L. and Ferriol, I. 2020. Detection of plant viruses and disease management: relevance of genetic diversity and evolution. *Front. Plant Sci.* 11:1092.
- Rubino, L. and Russo, M. 1998. Membrane targeting sequences in tombusvirus infections. *Virology* 252:431-437.
- Salesse-Smith, C. E., Sharwood, R. E., Busch, F. A., Kromdijk, J., Bardal, V. and Stern, D. B. 2018. Overexpression of Rubisco subunits with RAF1 increases Rubisco content in maize. *Nat. Plants* 4:802-810.
- Salonen, A., Ahola, T. and Kääriäinen, L. 2005. Viral RNA replication in association with cellular membranes. *Curr. Top. Microbiol. Immunol.* 285:139-173.
- Sanfaçon, H. 2005. Replication of positive-strand RNA viruses in plants: contact points between plant and virus components. *Can. J. Bot.* 83:1529-1549.
- Serrano, I., Audran, C. and Rivas, S. 2016. Chloroplasts at work during plant innate immunity. *J. Exp. Bot.* 67:3845-3854.
- Seyffert, C. and Tsuda, K. 2014. Salicylic acid signal transduction: the initiation of biosynthesis, perception and transcriptional reprogramming. *Front. Plant Sci.* 5:697.
- Shi, Y., Chen, J., Hong, X., Chen, J. and Adams, M. J. 2007. A potyvirus P1 protein interacts with the Rieske Fe/S protein of its host. *Mol Plant Pathol.* 8:785-790.
- Shimura, H., Pantaleo, V., Ishihara, T., Myojo, N., Inaba, J.-I., Sueda, K., Burgýán, J. and Masuta, C. 2011. A viral satellite RNA induces yellow symptoms on tobacco by targeting a gene involved in chlorophyll biosynthesis using the RNA silencing machinery. *PLoS Pathog.* 7:e1002021.
- Simon, A. E. and Miller, W. A. 2013. 3' cap-independent translation enhancers of plant viruses. *Annu. Rev. Microbiol.* 67:21-42.
- Stael, S., Kmiecik, P., Willems, P., Van der Kelen, K., Coll, N. S., Teige, M. and Van Breusegem, F. 2015. Plant innate immunity: sunny side up? *Trends Plant Sci.* 20:3-11.
- Thivierge, K., Cotton, S., Dufresne, P. J., Mathieu, I., Beauchemin, C., Ide, C., Fortin, M. G. and Laliberté, J.-F. 2008. Eukaryotic elongation factor 1A interacts with turnip mosaic virus RNA-dependent RNA polymerase and VPg-Pro in virus-induced vesicles. *Virology* 377:216-225.
- Torres, M. A., Jones, J. D. G. and Dangl, J. L. 2006. Reactive oxygen species signaling in response to pathogens. *Plant Physiol.* 141:373-378.
- Turner, J. G., Ellis, C. and Devoto, A. 2002. The jasmonate signal pathway. *Plant Cell* 14(Suppl):S153-S164.
- Wang, B., Li, Z., Ran, Q., Li, P., Peng, Z. and Zhang, J. 2018. *ZmNF-YB16* overexpression improves drought resistance and yield by enhancing photosynthesis and the antioxidant capacity of maize plants. *Front. Plant Sci.* 9:709.
- Wasternack, C. and Hause, B. 2013. Jasmonates: biosynthesis, perception, signal transduction and action in plant stress response, growth and development: an update to the 2007 review in Annals of Botany. *Ann. Bot.* 111:1021-1058.
- Wei, T., Zhang, C., Hou, X., Sanfaçon, H. and Wang, A. 2013. The SNARE protein Syp71 is essential for turnip mosaic virus infection by mediating fusion of virus-induced vesicles with chloroplasts. *PLoS Pathog.* 9:e1003378.
- Whitfield, A. E., Falk, B. W. and Rotenberg, D. 2015. Insect vector-mediated transmission of plant viruses. *Virology* 479-480:278-289.
- Widyasari, K., Alazem, M. and Kim, K.-H. 2020. Soybean resistance to soybean mosaic virus. *Plants* 9:219.
- Widyasari, K., Tran, P.-T., Shin, J., Son, H. and Kim, K.-H. 2022. Overexpression of purple acid phosphatase GmPAP2.1 confers resistance to soybean mosaic virus in a susceptible soybean cultivar. *J. Exp. Bot.* 73:1623-1642.
- Wildermuth, M. C., Dewdney, J., Wu, G. and Ausubel, F. M. 2001. Isochorismate synthase is required to synthesize salicylic acid for plant defense. *Nature* 414:562-565.

- Wu, J., Yang, R., Yang, Z., Yao, S., Zhao, S., Wang, Y., Li, P., Song, X., Jin, L., Zhou, T., Lan, Y., Xie, L., Zhou, X., Chu, C., Qi, Y., Cao, X. and Li, Y. 2017. ROS accumulation and antiviral defense control by microRNA528 in rice. *Nat. Plants* 3:16203.
- Xu, K. and Nagy, P. D. 2014. Expanding use of multi-origin subcellular membranes by positive-strand RNA viruses during replication. *Curr. Opin. Virol.* 9:119-126.
- Yang, F., Xiao, K., Pan, H. and Liu, J. 2021. Chloroplast: the emerging battlefield in plant-microbe interactions. *Front. Plant Sci.* 12:637853.
- Zhai, Y., Yuan, Q., Qiu, S., Li, S., Li, M., Zheng, H., Wu, G., Lu, Y., Peng, J., Rao, S., Chen, J. and Yan, F. 2021. Turnip mosaic virus impairs perinuclear chloroplast clustering to facilitate viral infection. *Plant Cell Environ.* 44:3681-3699.
- Zhang, C., Grosic, S., Whitham, S. A. and Hill, J. H. 2012. The requirement of multiple defense genes in soybean Rsv1-mediated extreme resistance to soybean mosaic virus. *Mol. Plant-Microbe Interact.* 25:1307-1313.
- Zhao, J., Liu, Q., Zhang, H., Jia, Q., Hong, Y. and Liu, Y. 2013. The RubisCO small subunit is involved in tobamovirus movement and Tm-2<sup>2</sup>-mediated extreme resistance. *Plant Physiol.* 161:374-383.
- Zhao, J., Xu, J., Chen, B., Cui, W., Zhou, Z., Song, X., Chen, Z., Zheng, H., Lin, L., Peng, J., Lu, Y., Deng, Z., Chen, J. and Yan, F. 2019. Characterization of proteins involved in chloroplast targeting disturbed by rice stripe virus by novel protoplast-chloroplast proteomics. *Int. J. Mol. Sci.* 20:253.
- Zhao, J., Zhang, X., Hong, Y. and Liu, Y. 2016. Chloroplast in plant-virus interaction. *Front. Microbiol.* 7:1565.
- Zurbriggen, M. D., Carrillo, N. and Hajirezaei, M.-R. 2010. ROS signaling in the hypersensitive response: when, where and what for? *Plant Signal. Behav.* 5:393-396.